#### TABLE VII

# PERCENTAGE EVALUATION FOR HEARING IMPAIRMENT (DIAGNOSTIC CODE 6100)

#### **Poorer Ear**

	XI	100*										
Better Ear	X	90	80									
	IX	80	70	60								
	VIII	70	60	50	50							
	VII	60	60	50	40	40						
	VI	50	50	40	40	30	30	And a common terminal transport				
Ř	V	40	40	40	30	30	20	20				
	IV	30	30	30	20	20	20	10	10			
	III	20	20	20	20	20	10	10	10	0		
	II	10	10	10	10	10	10	10	0 .	0	0	
-	I	10	10	0	0	0	0	0	0	. 0	0	0
i		XI	X	IX	VIII	VII	VI	V	IV	III	11	I

<sup>\*</sup> Review for entitlement to special monthly compensation under §3.350 of this chapter.

[64 FR 25206, May 11, 1999]

## § 4.86 Exceptional patterns of hearing impairment.

(a) When the puretone threshold at each of the four specified frequencies (1000, 2000, 3000, and 4000 Hertz) is 55 decibels or more, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. Each ear will be evaluated separately.

(b) When the puretone threshold is 30 decibels or less at 1000 Hertz, and 70 decibels or more at 2000 Hertz, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. That numeral will then be elevated to the next higher

#### **Department of Veterans Affairs**

§ 4.88a

Roman numeral. Each ear will be evaluated separately.

(Authority: 38 U.S.C. 1155) [64 FR 25209, May 11, 1999]

#### §4.87 Schedule of ratings—ear.

#### DISEASES OF THE EAR

DISEASES OF THE EAR					
6200 Chronic suppurative otitis media, mastoiditis, or cholesteatoma (or any combination):  During suppuration, or with aural polyps  NOTE: Evaluate hearing impairment, and complications such as labyrinthitis, tinnitus, facial nerve paralysis, or bone loss of skull, separately.					
factory.  6201 Chronic nonsuppurative otitis media with effusion (serous otitis media): Rate hearing impairment 6202 Otosclerosis: Rate hearing impairment 6204 Peripheral vestibular disorders: Dizziness and occasional staggering					
Occasional dizziness					
6205 Meniere's syndrome (endolymphatic hydrops): Hearing impairment with attacks of vertigo and cerebellar gait occurring more than once weekly, with or without tinnitus Hearing impairment with attacks of vertigo and cerebellar gait occurring from one to four times a month, with or without tinnitus					
Complete loss of both Complete loss of one Deformity of one, with loss of one-third or more of the substance 6208 Malignant neoplasm of the ear (other than skin only) NOTE: A rating of 100 percent shall continue be- yond the cessation of any surgical, radiation treatment, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appro- priate disability rating shall be determined by mandatory VA examination. Any change in evaluation based on that or any subsequent					

examination shall be subject to the provisions

of §3.105(e) of this chapter. If there has been

no local recurrence or metastasis, rate on re-

6209 Benign neoplasms of the ear (other than skin

Rate on impairment of function.

6210 Chronic otitis externa:

DISEASES OF THE EAR—Continued

	Rat- ing
Swelling, dry and scaly or serous discharge, and itching requiring frequent and prolonged treatment	10 0 10

(Authority: 38 U.S.C. 1155) [64 FR 25210, May 11, 1999]

Rating

10

30 10

100

60

30

### § 4.87a Schedule of ratings—other sense organs.

	ing
6275 Sense of smell, complete loss	10
6276 Sense of taste, complete loss	10
NOTE: Evaluation will be assigned under diag- nostic codes 6275 or 6276 only if there is an anatomical or pathological basis for the condi- tion.	

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999]

INFECTIOUS DISEASES, IMMUNE DIS-ORDERS AND NUTRITIONAL DEFI-CIENCIES

#### §4.88 [Reserved]

#### §4.88a Chronic fatigue syndrome.

- (a) For VA purposes, the diagnosis of chronic fatigue syndrome requires:
- (1) new onset of debilitating fatigue severe enough to reduce daily activity to less than 50 percent of the usual level for at least six months; and
- (2) the exclusion, by history, physical examination, and laboratory tests, of all other clinical conditions that may produce similar symptoms; and
  - (3) six or more of the following:
  - (i) acute onset of the condition,
  - (ii) low grade fever,
  - (iii) nonexudative pharyngitis,
- (iv) palpable or tender cervical or axillary lymph nodes,
- (v) generalized muscle aches or weakness,
- (vi) fatigue lasting 24 hours or longer after exercise,